

A Landau's theorem in several complex variables

Cinzia Bisi¹

Received: 28 January 2016 / Accepted: 2 July 2016 / Published online: 29 July 2016
© Fondazione Annali di Matematica Pura ed Applicata and Springer-Verlag Berlin Heidelberg 2016

Abstract In one complex variable, it is well known that if we consider the family of all holomorphic functions on the unit disc that fix the origin and with first derivative equal to 1 at the origin, then there exists a constant ρ , independent of the functions, such that in the image of the unit disc of any of the functions of the family, there is a disc of universal radius ρ . This is the so celebrated Landau's theorem. Many counterexamples to an analogous result in several complex variables exist. In this paper, we introduce a class of holomorphic maps for which one can get a Landau's theorem and a Brody–Zalcman theorem in several complex variables.

Keywords Normal family · Brody–Zalcman theorem · Landau's constant · Rouché's theorem

Mathematics Subject Classification 32H99 · 32A18 · 30D45

1 Introduction

An important tool in geometric function theory in one complex variable is the so-called Landau's theorem.

It says: *if we consider the family \mathcal{F} , of holomorphic functions in the unit disc Δ of \mathbb{C} such that $f(0) = 0$, $f'(0) = 1$, then there is a constant $\rho > 0$, independent of f , such that $f(\Delta)$ contains a disc of radius ρ [8, 16–18].*

Partially supported by GNSAGA of the INdAM, by FIRB “Geometria Differenziale e teoria geometrica delle funzioni” and by PRIN “Varietà reali e complesse: geometria, topologia ed analisi armonica”.

✉ Cinzia Bisi
bsicnz@unife.it

¹ Dipartimento di Matematica e Informatica, Università degli Studi di Ferrara,
Via Machiavelli 35, 44121 Ferrara, Italy

By the *Landau's number* $l(f)$ of f is meant the supremum of the set of positive numbers r such that $f(\Delta)$ contains a disc of radius r . By the *Landau's constant* ρ , we meant $\inf_{f \in \mathcal{F}} l(f)$.

An easy consequence of such result is the Picard theorem: *a non-constant entire function $f: \mathbb{C} \rightarrow \mathbb{C}$ can omit at most one point.*

A related result is the compactness result of Brody–Zalcman: *if \mathcal{H} is a non-relative compact family of holomorphic maps in the unit disc, then after reparametrization we can get an extracted subsequence of functions in \mathcal{H} converging to a non-constant entire map [6, 25, 26].*

It is well known that such principles break down in several complex variables in a spectacular way.

There are holomorphic maps $F: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ with Jacobian equal to 1, and whose image omits a non-empty open set. These maps play an important role in holomorphic dynamics, see [22].

To be more explicit, let $h(z, w) = (z^2 + aw, z)$, with $|a| < 1$, be a so-called h enon map of \mathbb{C}^2 : it fixes $(0, 0)$ and the attraction domain Ω of $(0, 0)$ is a Fatou–Bieberbach domain isomorphic to \mathbb{C}^2 . Denote with $\Psi_h: \mathbb{C}^2 \rightarrow \Omega$ such isomorphism such that $\Psi'_h(0, 0) = \text{Id}$. Let \bar{h} be the meromorphic extension of h to \mathbb{P}^2 . If (z, w) are the coordinates of \mathbb{C}^2 and $[z:w:t]$ are the coordinates of \mathbb{P}^2 , then the line at infinity has equation $\{t = 0\}$. We denote, respectively, with I^+ and I^- the indeterminacy locus of \bar{h} and h^{-1} ; they are two isolated points in $\{t = 0\}$. It is well known, see [22], that the h enon map h has $I^- = [1:0:0]$, and it is an attracting fixed point at the line at infinity. Then, Ω omits the basin of attraction of I^- , which is an open set. Hence, it cannot be that each $\Psi_{h,n}(z, w) := \frac{1}{n}\Psi_h(nz, nw)$ is such that the image of $B(0, 1)$ contains a ball $B(0, r)$, because if this is the case, then $\Psi_h(nB(0, 1)) \supset nB(0, r)$ and finally $\Psi_h(\mathbb{C}^2) \supset \mathbb{C}^2$, which is not true because Ψ_h omits an open set.

There are also some elder counterexamples, one by L.A. Harris, published in 1977 [15] that we recall for the sake of completeness: given $\delta > 0$, choose n so that $n\delta^2 > 2$ and denote $g(z, w) = (z + nw^2, w)$. Suppose that the image of the open polydisc by g contains a ball of radius δ and let (α_0, β_0) be its center. Then, given $|\zeta| < \delta$, there exist points (z_0, w_0) and (z_1, w_1) in the polydisc such that $g(z_0, w_0) = (\alpha_0, \beta_0)$ and $g(z_1, w_1) = (\alpha_0, \beta_0 + \zeta)$. Hence, $w_1 - w_0 = \zeta, w_1 + w_0 = 2\beta_0 + \zeta$, and $n(w_1^2 - w_0^2) = z_0 - z_1$, so $n|\zeta||2\beta_0 + \zeta| \leq 2$, for all $|\zeta| < \delta$. Thus, $n\delta^2 \leq 2$, the desired contradiction.

In the same spirit, a second counterexample was given by Duren and Rudin [9]: if $\delta > 0$, then the map $f(z, w) = (z, w + (\frac{z}{\delta})^2)$ is in the class of all biholomorphic maps from the unit polydisc into \mathbb{C}^2 which fix the origin and whose Jacobian matrix is the identity at the origin, but the image under f of the polydisc contains no closed ball of radius δ .

Indeed, for no $(u, v) \in \mathbb{C}^2$, the image by f of the polydisc Δ^2 contains the circle:

$$C = \left\{ (u + \delta e^{i\theta}, v): -\pi \leq \theta \leq \pi \right\}.$$

To see this, fix $(u, v) \in \mathbb{C}^2$. If $(u + \lambda, v) \in f(\Delta^2)$ then, by definition of f , we have that:

$$|v - \delta^{-2}(u + \lambda)^2| < 1.$$

Therefore, if all points of C were in $f(\Delta^2)$, the inequality:

$$|(\delta^2 v - u^2) - 2u\delta e^{i\theta} - \delta^2 e^{2i\theta}| < \delta^2$$

would hold for all θ . Parseval's equality shows that this is impossible.

Even if several weak versions of the Landau's theorem in several complex variables have already been given, see, for instance, [7, 10, 12, 13, 19, 23, 24]; nevertheless, the author believes that this paper can add something to the already existing literature: indeed, the purpose of this

note is to introduce a class of holomorphic maps for which one can get a Landau’s theorem and a Brody–Zalcman theorem in more than one variable and to underline the connection among the two.

Recently, the Landau’s theorem has also bring the attention of people working over the quaternionic variable, see, for instance, [1–5].

2 A theorem of Brody–Zalcman type

Let $\Phi: \mathbb{B}^k \rightarrow \mathbb{C}^k$ where \mathbb{B}^k is the unit ball of \mathbb{C}^k , $k > 1$. Let $\Phi'(a)$ denote the Jacobian matrix of Φ computed in the point a .

Theorem 2.1 (Brody–Zalcman-type Theorem) *Let C be a positive constant. Consider a family of such holomorphic maps Φ satisfying*

$$\|\Phi'(a)\| \cdot \|\Phi'(a)^{-1}\| \leq C, \quad \forall a \in \mathbb{B}^k. \tag{1}$$

If the family is not normal, then, after reparametrization, we can extract a subsequence converging to a non-degenerate holomorphic map $\Psi: \mathbb{C}^k \rightarrow \mathbb{C}^k$.

Proof Let Φ_n be an arbitrary sequence of maps of the family. We can assume that the maps Φ_n are defined in a neighborhood of $\overline{\mathbb{B}^k}$.

Define

$$\lambda_n := \sup_{|z|<1} (1 - |z|) \|\Phi'_n(z)\|.$$

We can also assume that $\lambda_n \rightarrow +\infty$, because if not the family is normal [21]. Let a_n be such that $(1 - |a_n|) \|\Phi'_n(a_n)\| = \lambda_n$.

Define

$$B_n := [(\Phi'_n)(a_n)]^{-1}$$

and

$$\Psi_n(z) := \Phi_n(a_n + B_n z).$$

We are going to show that $\Psi'_n(z)$ is well defined in $|z| \leq \frac{\lambda_n}{2C}$.

Indeed, $\Psi'_n(z) = \Phi'_n(a_n + B_n z) \circ B_n$ with $\Psi'_n(0) = \text{Id}$ and since

$$(1 - |a_n + B_n z|) \|\Phi'_n(a_n + B_n z)\| \leq \lambda_n$$

we have:

$$\|\Psi'_n(z)\| \leq \|\Phi'_n(a_n + B_n z)\| \cdot \|B_n\| \leq \frac{\lambda_n \|B_n\|}{1 - |a_n| - |B_n z|}. \tag{2}$$

If $|z| \leq \frac{\lambda_n}{2C}$, then by (1):

$$|B_n z| \leq \frac{C|z|}{\|\Phi'(a_n)\|} \leq \frac{(1 - |a_n|)}{\lambda_n} C \frac{\lambda_n}{2C} = \frac{1 - |a_n|}{2}. \tag{3}$$

So

$$\|\Psi'_n(z)\| \leq \frac{2\lambda_n}{1 - |a_n|} \|B_n\| \leq \frac{2\lambda_n}{1 - |a_n|} \frac{C}{\|\Phi'_n(a_n)\|} \leq 2C$$

So the family Ψ_n is locally normal in $|z| \leq \frac{\lambda_n}{C} \rightarrow \infty$.

We get that Ψ_n tends to a holomorphic map $\mathbb{C}^k \rightarrow \mathbb{C}^k$. Moreover, $\Psi'_n(0) = \text{Id}$ so $\Psi'(0) = \text{Id}$. Hence, Ψ is non-degenerate. □

Remark 2.2 If $k = 1$, then (1) is automatically satisfied.

Remark 2.3 Condition (1) implies that all the eigenvalues are comparable, i.e., $|\lambda_{\max}(\Phi'(a))| \leq C|\lambda_{\min}(\Phi'(a))|$ for all $a \in \mathbb{B}^k$, where $|\lambda_{\min}(\Phi'(a))|$ and $|\lambda_{\max}(\Phi'(a))|$ are, respectively, the minimal and the maximal modulus of the eigenvalues of the Jacobian matrix $\Phi'(a)$.

Remark 2.4 Furthermore, it is enough to assume (1) out of an analytic set, so the maps Φ do not need to be locally invertible. Indeed, if we suppose that (1) holds out of an analytic set A_Φ , we can choose $a_n \notin A_{\Phi_n}$.

Remark 2.5 The condition (1) can be refined in

$$\sup_{|z| \leq \frac{1-|a|}{2}} \|\Phi'(a+z) \cdot \Phi'(a)^{-1}\| \leq C$$

This condition is less strong of (1) and implies that $\|\Psi'_n\|$ is bounded for $|z| \leq d_n$ with $d_n \rightarrow +\infty$.

Remark 2.6 The Brody–Zalcman renormalization Theorem 2.1 works also for families of maps from \mathbb{B}^k to a compact Hermitian manifold M of dimension k [11].

We also point out that, in [20], several sufficient conditions for a family of quasiregular mappings to be normal were previously given.

3 Landau’s theorem

Assume that $\Phi: \mathbb{B}^k \rightarrow \mathbb{C}^k$, $\Phi(0) = 0$, and $\Phi'(0) = \text{Id}$.

Theorem 3.1 (Landau’s theorem) *Let C be a positive constant. Assume*

$$\|\Phi'(z)\| \cdot \|\Phi'(z)^{-1}\| \leq C, \quad \forall z \in \mathbb{B}^k, \tag{4}$$

then there exists $\rho > 0$, depending only on C , such that $\Phi(\mathbb{B}^k)$ contains a ball of radius $\rho > 0$. There exists also a domain U such that $\Phi(U) = B(a, \rho)$.

Proof Suppose, by contradiction, that this is not the case. Then, there exists a sequence Φ_n such that $\Phi_n(0) = 0$, $\Phi'_n(0) = \text{Id}$ satisfying (4) and not the conclusion of the theorem.

If $(1-|z|)\|\Phi'_n(z)\| \leq A$, with A constant, then Φ_n is normal, and we can assume $\Phi_n \rightarrow \Psi$, with $\Psi(0) = 0$ and $\Psi'(0) = \text{Id}$.

On an appropriate sphere $\partial B(0, r)$, we get $|\Phi_n - \Psi| < |\Psi|$. So by Rouché’s theorem and by the fact that the image of Ψ contains a ball of radius R , we get that eventually also the images of Φ_n contain a ball of radius smaller or equal than R because we are interested in the inequality

$$|\Phi_n - b - \Psi + b| < |\Psi - b| \leq |\Psi| + |b|,$$

for all $b \in B(a, R)$ and this will be satisfied possibly on a $\partial B(0, r')$ with $r' \leq r$ which is a contradiction.

Hence, we can assume that $\sup_{|z|<1} (1 - |z|)|\Phi'_n(z)| = \lambda_n \rightarrow +\infty$.
 As in the previous result, we can reparametrize and get

$$\Psi_n := \Phi_n(a_n + A_n z) \rightarrow \Psi, \quad \Psi'(0) = \text{Id}.$$

Then, $\Psi(\mathbb{B}^k)$ contains a ball centered at $\Psi(0) = a$, $B(\Psi(0), R)$.

Since by Rouché’s theorem:

$$|\Psi_n - \Psi| < |\Psi|$$

on $\partial B(0, r)$, then eventually the image of Ψ_n contains a ball of radius smaller or equal than R because we are interested in the inequality

$$|\Psi_n - b - \Psi + b| < |\Psi - b| \leq |\Psi| + |b|,$$

for all $b \in B(a, R)$ and this will be satisfied possibly on a $\partial B(0, r')$ with $r' \leq r$.

Then, the image of Φ_n will contain eventually a ball of radius $R' \leq R$ and we get the domain U .

The supremum of the set of positive numbers R' such that $\Phi(\mathbb{B}^k)$ contains a ball of radius R' is the so-called *Landau’s number* $l(\Phi)$. The constant $\rho = \inf_{\Phi \in \mathcal{G}} l(\Phi)$ where \mathcal{G} is the set of $\Phi: \mathbb{B}^k \rightarrow \mathbb{C}^k$, $\Phi(0) = 0$, and $\Phi'(0) = \text{Id}$ satisfying condition (4). □

We point out that, following the same arguments of [14], it is possible to find an open set U on which the entire family of Φ ’s satisfying (4) is injective, and it is also possible to relate the constant C with the uniform radius ρ .

Example 3.2 Consider the following family of hénon maps of \mathbb{C}^2 :

$$h_b(z, w) = (z^2 + bw, z)$$

with $\delta < |b| \leq \eta$, for δ, η fixed constants.

Let $f(z, w) = (e^{cz} - 1, e^{cw} - 1)$, with $|c|$ small enough if you need f invertible, then the family

$$g_{b,c}(z, w) = \{h_b \circ (e^{cz} - 1, e^{cw} - 1)\}$$

satisfies the hypothesis of Theorem 3.1, up to dilation.

Furthermore, if the ball centered in the origin is enough small, then the ball of universal radius contained in the images of the family is a ball on which the maps of the family are invertible.

Corollary 3.3 *Let $f: \mathbb{C}^k \rightarrow \mathbb{C}^k$ be an entire map. Suppose that $f'(0) = \text{Id}$ and suppose that there exists a constant C such that $\|f'(z)\| \cdot \|f'(z)^{-1}\| \leq C$, then $f(\mathbb{C}^k)$ contains balls of arbitrary large radius.*

Proof Indeed, $\frac{1}{R} f(R \cdot B(0, 1)) \supset B(a, r)$ for any $R > 0$, by Theorem 3.1. □

Acknowledgements The author warmly thanks the Referee for his/her accurate remarks.

References

1. Bisi, C., Gentili, G.: Möbius transformations and the Poincaré distance in the quaternionic setting. Indiana Univ. Math. J. **58**(6), 2729–2764 (2009)

2. Bisi, C., Gentili, G.: On the geometry of the quaternionic unit disc. In: Sabadini, I., Sommen, F. (eds.) *Hypercomplex analysis and applications*. Trends in Mathematics, pp. 1–11. Springer, Basel (2011)
3. Bisi, C., Stoppato, C.: The Schwarz–Pick lemma for slice regular functions. *Indiana Univ. Math. J* **61**(1), 297–317 (2012)
4. Bisi, C., Stoppato, C.: Regular vs. classical Möbius transformations of the quaternionic unit ball. In: Gentili, G., Sabadini, I., Shapiro, M., Sommen, F., Struppa, D.C. (eds.) *Advances in hypercomplex analysis*. Springer INdAM Series, vol. 1, pp. 1–13. Springer, Milan (2013)
5. Bisi, C., Stoppato, C.: Landau’s theorem for slice regular functions on the quaternionic unit ball, Preprint (2016)
6. Brody, R.: Compact manifolds and hyperbolicity. *Trans. Am. Math. Soc.* **235**, 213–219 (1978)
7. Chen, H., Gauthier, P.M.: Bloch constants in several variables. *Trans. Am. Math. Soc* **353**(4), 1371–1386 (2001)
8. Conway, J.B.: *Functions of One Complex Variable*. Graduate Texts in Mathematics, vol. 11. Springer, New York (1973)
9. Duren, P., Rudin, W.: Distortion in several variables. *Complex Var. Theory Appl.* **5**(2–4), 323–326 (1986)
10. FitzGerald, C.H., Gong, S.: The Bloch theorem in several complex variables. *J. Geom. Anal.* **4**(1), 35–58 (1994)
11. Fornæss, J.E., Stout, E.L.: Regular holomorphic images of balls. *Ann. Inst. Fourier* **32**(2), 23–36 (1982)
12. Graham, I., Kohr, G.: *Geometric function Theory in One and Higher Dimensions*. Pure and Applied Mathematics, vol. 255. Dekker, New York (2003)
13. Graham, I., Varolin, D.: Bloch constants in one and several variables. *Pac. J. Math.* **174**(2), 347–357 (1996)
14. Hahn, K.T.: Higher dimensional generalizations of the Bloch constant and their lower bounds. *Trans. Am. Math. Soc.* **179**, 263–274 (1973)
15. Harris, L.A.: On the size of balls covered by analytic transformations. *Mon. Math.* **83**, 9–23 (1977)
16. Heins M.: *Selected topics in the classical theory of functions of a complex variable*. Athena Series: Selected Topics in Mathematics Holt, Rinehart and Winston, New York, (1962)
17. Landau, E.: Über die Blochsche Konstante und zwei verwandte Weltkonstanten. *Math. Z* **30**(1), 608–634 (1929)
18. Landau, E.: *Darstellung und Begründung einiger neuerer Ergebnisse der Funktionentheorie*. Springer, Berlin (1929)
19. Liu, X.Y.: Bloch functions of several complex variables. *Pac. J. Math.* **152**(2), 347–363 (1992)
20. Miniowitz, R.: Normal families of quasimeromorphic mappings. *Proc. Am. Math. Soc.* **84**(1), 35–43 (1982)
21. Schiff, J.L.: *Normal families*. Universitext. Springer, New York (1993)
22. Sibony, N.: Dynamique des applications rationnelles de \mathbb{P}^k . (French) [Dynamics of rational maps of \mathbb{P}^k] *Dynamique et géométrie complexes* (Lyon, 1997), Panor. Synthèses, vol. 8, Soc. Math. France, Paris (1999), pp. 97–185 (1997)
23. Takahashi, S.: Univalent mappings in several complex variables. *Ann. Math.* **53**(2), 464–471 (1951)
24. Wu, H.: Normal families of holomorphic mappings. *Acta Math.* **119**, 193–233 (1967)
25. Zalcman, L.: A heuristic principle in complex function. *Am. Math. Mon.* **82**(8), 813–818 (1975)
26. Zalcman, L.: Normal families: new perspectives. *Bull. (New Series) Am. Math. Soc.* **35**(3), 215–230 (1998)