


Gli APTAMERI

APTAMERI


Acidi nucleici a singolo filamento caratterizzati da una specifica **struttura tridimensionale** che si lega direttamente alla proteina target.



Interazione Acido Nucleico/Proteina


Aptameri

Dimensioni: 30-70 nucleotidi




Folding

Molecola Lineare




Struttura
tridimensionale
stabile




Anatomia degli Aptameri


Gli aptameri sono molecole selezionate per legarsi
in modo specifico ad una predefinita *proteina*
target



Selezione in vitro degli Aptameri:

SELEX (systematic evolution of ligands by exponential enrichment)







Selezione in vitro degli Aptameri:

SELEX (systematic evolution of ligands by exponential enrichment)

1. Sintesi chimica di 10^{14} RNA o DNA (Libreria)
2. Incubazione con la proteina target: chromatografia per affinità
3. Rimozione degli oligo *non legati* mediante buffer di lavaggio
4. Rimozione degli oligo *legati* alla proteina target con una soluzione contenente la proteina target
5. Retrotrascrizione e PCR (RNA) o solo PCR (DNA) degli oligo che si sono legati
6. Trascrizione in vitro (RNA) o solo denaturazione (DNA) per separare i filamenti
7. Inizio di un nuovo ciclo fino a 5-10 cicli





Applicazioni degli Aptameri:

- 1. Ricerca**
- 2. Diagnostica**
- 3. Terapia**

Applicazioni degli Aptameri:

ALTERNATIVA AGLI ANTICORPI

- elevate specificità e affinità unite a ridotte dimensioni
- sintesi chimica (vs sintesi in animali o colture cellulari)
- facilmente modificabili: marcatura con radioattivo, code fluorescenti e biotinilate...

APPLICAZIONI IN VIVO:

- nessuna tossicità dimostrata (facilmente eliminabili da sangue e reni)
- non immunogenici
- tessuto-specifici

Table 3
Examples of applied aptamers

Target	Aptamer/assay	Field of application	Research or product state	Reference
VEGF	RNA aptamer, chemically modified DNA aptamer (thrombin inhibitor ARC-138)	Therapy, wet age related macula degeneration	Product: Macugen® (Pfizer Inc.)	Maberley (2005) and Chapman and Beckey (2006)
Thrombin		Therapy, anticoagulant	Product development, Phase 1 studies in August 2004 (Archemix Corp.)	Nimjee et al. (2005a)
Factor IXa	RNA aptamer (factor IXa inhibitor) and its antidote REGI	Therapy, anticoagulant	Product development, Phase 1 studies completed in 2006, (Regado Biosciences Inc.; Archemix Corp.)	Nimjee et al. (2005a) and Dyke et al. (2006)

TERAPIA:

Condizioni patologiche acute e spazialmente confinate

→ **Trombosi: aptameri contro trombina, FVIIa, FIXa**

→ **Cancro: aptameri contro proteine segnale
(es. Crescita, differenziazione,
trasformazione cellulare...)**

→ **Patologie virali: identificazione e inibizione
di proteine virali**

Gli APTAMERI: un'applicazione

Rusconi CP, Scardino E, Layzer J, Pitoc GA,
Ortel TL, Monroe D, Sullenger BA

**RNA aptamers as reversible
antagonists of coagulation
factor IXa**

Nature 2002; 419: 90-94
[\(\[www.nature.com\]\(http://www.nature.com\)\)](http://www.nature.com)

COAGULAZIONE DEL SANGUE

Danno vascolare




Attivazione a cascata di fattori e
cofattori plasmatici




Complessi macromolecolari


Substrato
inattivo



Complesso di attivazione del FX



Complesso di attivazione del FX




- {
1. Elevata AFFINITA' con il FIXa
 2. SPECIFICITA' per il FIXa

Selezione degli Aptameri

SELEX

Libreria: 10^{14}
oligonucleotidi (RNA)

Retrotrascrizione degli
RNA selezionati e
sequenziamento




1. AFFINITA' con il FIXa

→

Aptamer	S1	L1	S2	L2	S2	L3	S1	
9-3	5' <u>GGGAUGGGGA</u>	CUAUACC	GCG	UAAAUGC	UGC	C	UCCCCAUUCC	GGNACGCU 3'
9-20	5' <u>GGGCA</u>	CUAUACCG	GCA	AUCG	UGC	A	UCCC	UGGACCUAACAAUA 3'
9-19	5' <u>GGaUGGGA</u>	CCAUUA	AGGA	CUAC	UCGU	GAA	UCCGACCC	ADCAAGGCGCACAA 3'
9-4	5' <u>GGGaUGGGC</u>	ACUAUAC	GCA	UCU	UGC	U	GGCUGGCC	GGGAGUCAAUUG 3'
9-12	5' <u>GGGAUGGG</u>	CGAUUA	UAC	ACATUG	GUG	AU	CCCACCC	ACAUGAAAACCACAG 3'
9-17	5' <u>GAGGqauGGGA</u>	CCAUAC	GCA	CAU	UGC	UGAA	UCCCGCCUC	AAUAGCACCACUAC 3'
9-25	5' <u>GGGAUGGGGA</u>	CCAUUA	ACTC	UAAC	GGGU	GAA	UCCCGCAUCUC	QACAAUA 3'
9-26	5' <u>GGGAUGGG</u>	UGAUUA	ACCA	CUC	UGGU	GAA	CCCCeUCCC	GACUUGCGCGCA 3'
9-11	5' <u>GGGauUGGGA</u>	CUAUUA	UUCCG	AAU	CUGGA	C	UCCCGACCU	GGCUGCCCCAGA 3'
9-2	5' <u>GGGAUGGG</u>	CUAUUA	CAC	GCUG	GUG	AU	CCCACUC	AAUUGAAAACAAUA 3'
9-7	5' <u>GGGAUGGG</u>	CGAUUA	ACCA	ACA	UGGU	GAU	CCCCUUC	ADCAUACCCUACAA 3'
9-28	5' <u>GGGAUGGGCG</u>	CCAUAC	GCA	CAU	UGC	UGCAU	GGCCUUCUCCC	GUAGAAC 3'
9-16	5' <u>QAGGGaUGGG</u>	CCAUAC	GUCC	ACGA	CUQC	A	CCCGaCCCUU	CAGCCCCAGGUC 3'
9-18	5' <u>GGGAUGGG</u>	CCAUUA	ACCA	CUU	UGGU	GAA	CCCACCC	AGCUCCUGUGAUUG 3'
9-14	5' <u>GGGAUGGGGA</u>	CUAUUA	OGU	GAACG	ACU	GCA	UCCCaCUUCCC	GGCCAUUGC 3'
9-27	5' <u>GGGaUGGG</u>	UAAAUA	ACU	GUU	UGG	UGAA	CCCCACCC	AAACUCCCCAUUCCUA 3'

Aptamero
9.3t




9.3 tM:
controllo
negativo
(aptamero
inattivo)

2. SPECIFICITA' per il FIXa

Aptamero
9.3t

5000 volte più
specifico



FVIIa

FXa

FXIa

APC

Serin-proteasi

Inibizione IN VITRO dell'attività del FIXa


Assemblamento del complesso



Introduzione dell'aptamero 9.3t



Inibizione IN VITRO dell'attività del FIXa



9.3t: aptamero selezionato
9.3tM: controllo negativo


L'aptamero **blocca** l'attività del **FIXa**
in vitro

Inibizione in plasma umano dell'attività del FIXa

Procedimento:

1. Aggiunta di diverse concentrazioni di aptamero (9.3t) e controllo negativo (9.3tM) a plasma umano
2. Misurazione del tempo di coagulazione del plasma

Inibizione IN VIVO dell'attività del FIXa




9.3t prolunga il tempo di coagulazione in modo dose-dipendente

L'aptamero inibisce l'attività del FIXa
in vivo

Reversibilità dell'azione dell'aptamero: ANTIDOTO

Antidoto = oligo complementare all'aptamero, in grado di alterare la sua conformazione




Reversibilità dell'azione dell'aptamero: ANTIDOTO

Plasma + aptamero = inibizione della coagulazione




Plasma non coagulato + antidoto = coagulazione




**Si misura la capacità del plasma di coagulare
entro 10 min.**

Reversibilità dell'azione dell'aptamero: ANTIDOTO



L'antidoto neutralizza l'azione
dell'aptamero


Reversibilità dell'azione dell'aptamero: ANTIDOTO



L'azione dell'antidoto è rapida e dose-dipendente


Efficacia di aptamero e antidoto su pazienti

6 Pazienti con trombosi → pazienti non sottoponibili
ai normali trattamenti anticoagulanti





First-in-human experience of an antidote-controlled anticoagulant using RNA aptamer technology: a phase 1a pharmacodynamic evaluation of a drug-antidote pair for the controlled regulation of factor IXa activity.

- Selectivity, titratability, rapidity of onset, and active reversibility are desirable pharmacological properties of anticoagulant therapy administered for acute indications and collectively represent an attractive platform to maximize patient safety. A novel anticoagulation system (REG1, Regado Biosciences), developed using a protein-binding oligonucleotide to factor IXa (drug, RB006) and its complementary oligonucleotide antidote (RB007), was evaluated in healthy volunteers. The primary objective was to determine the safety profile and to characterize the pharmacodynamic responses in this first-in-human study. METHODS AND RESULTS: Regado 1a was a subject-blinded, dose-escalation, placebo-controlled study that randomized 85 healthy volunteers to receive a bolus of drug or placebo followed 3 hours later by a bolus of antidote or placebo. Pharmacodynamic samples were collected serially. Subject characteristics were the following: median age, 32 years (interquartile range, 23 to 39 years); female gender, 35%; and median weight, 79 kg (interquartile range, 70 to 87 kg). No significant differences were found in median hemoglobin, platelet, creatinine, or liver function studies. There were no significant bleeding signals associated with RB006, and overall, both drug and antidote were well tolerated. One serious adverse event, an episode of transient encephalopathy, occurred in a subject receiving the low intermediate dose of RB006. The subject's symptoms resolved rapidly, and no further sequelae occurred. A predictable dose-pharmacodynamic response, reflected in activated partial thromboplastin time measurements, was seen after administration of the bolus of drug, with a clear correlation between the peak posttreatment activated partial thromboplastin time and post hoc weight-adjusted dose of drug (correlation coefficient, 0.725; $P < 0.001$). In subjects treated with drug, antidote administration reversed the pharmacological activity of the drug, with a rapid (mean time, 1 to 5 minutes across all dose levels) and sustained return of activated partial thromboplastin time to within the normal range. The activated clotting time followed a similar anticoagulant response and reversal pattern. As anticipated, prothrombin time remained unchanged compared with baseline. CONCLUSIONS: These observations represent a first-in-human experience of an RNA aptamer and its complementary oligonucleotide antidote used as an anticoagulant system. The findings contribute to an emerging platform of selective, actively reversible anticoagulant drugs for use among patients with thrombotic disorders of the venous and arterial circulations.



P polyethylene glycol; idT, inverted deoxythymidine





Pharmacodynamic effects of RB006 at 0 to 3 hours after RB006 administration.

The relative increase in APTT over baseline for each subject receiving RB006 before RB007 or placebo administration (all subjects assigned to arms 2 and 3) is shown vs subjects receiving placebo. Data represents the mean SEM for all subjects receiving treatment at each dose level.

CONCLUSIONI

- 1. Aptameri contro il FIXa sono potenti anticoagulanti**
- 2. Oligonucleotidi complementari agli aptameri possono agire da antidoti e neutralizzare l'azione anticoagulante**

Vascular Endothelial Growth Factor and the Potential Therapeutic Use of Pegaptanib (Macugen®) in Diabetic Retinopathy


Starita C, Patel M, Katz B, Adamis A

- Both clinical and preclinical findings have implicated vascular endothelial growth factor (VEGF) in the pathophysiology macular edema and degeneration.
- VEGF is both a potent enhancer of vascular permeability and a key inducer of angiogenesis.
- VEGF levels are elevated in the eyes of patients. Moreover, injection of VEGF (the VEGF165 isoform in particular) into healthy eyes of animals can induce associated ocular pathologies.
- Pegaptanib, a novel RNA aptamer currently used in the treatment of age-related macular degeneration, binds and inactivates VEGF165 and has been shown in animal models to reverse the blood-retinal barrier breakdown.


Il pegaptanib e` un antagonista selettivo del VEGF₁₆₅

Il Pegaptanib e` un aptamero a filamento singolo di **RNA** formato da 28 nucleotidi legato a 2 molecole di 20-kDa di glicole polietilenico (PEG)


E` dotato di alta affinita` per il VEGF₁₆₅ (**vascolarizzazione patologica**) e nessun legame con il VEGF₁₂₁ (**vascolarizzazione fisiologica**)





a | Sequenza e struttura secondaria del pegaptanib.




Il legame avviene tra la cisteina – 137 del VEGF₁₆₅ e l'uridina-14 dell'aptamero₁₄ (in rosso).








Extracellular Neutralization of VEGF



- Il Pegaptanib si lega specificamente al VEGF-165, impedendone l'aggancio con il suo recettore

Mantenimento o miglioramento del visus




*Nominal $P<0.05$ vs Usual Care

**Prespecified endpoint

Gragoudas ES et al. *N Eng J Med*. 2004;351:2805-2816.

Responders nello studio V.I.S.I.O.N. Lesioni iniziali vs tutti i Pazienti

- * Lesione iniziale definita come: occulta, senza essudati, e occhio controlaterale con visus migliore



V.I.S.I.O.N. Clinical Trial Group. *Retina* 2005;25:815-827.

Varie

What

Latin: “aptus” – to fit

Any protein or nucleic acid (RNA and DNA) sequence discovered through in vitro selection (SELEX) to recognize and specifically bind a target of interest


Sequences are selected from very large pools (10^{15} or greater)

Selection occurs using affinity chromatography

Non-bound sequences are eluted using several column volumes of buffer

Bound sequences are eluted with a solution containing the target compound

EQUILIBRIO EMOSTATICO




Sommario:

1. Definizione di Aptamero
2. Anatomia degli Aptameri
3. Selezione in vitro degli Aptameri
4. Un'applicazione: aptamero contro il FIXa
5. Altre Applicazioni

ssDNA separation methods

(-)primer modification: 3'-riboU




follow-up treatment: → alkaline hydrolysis
(selective disruption of the (-)strand at the riboU-position)



truncation of 18 nucleotides from the (-)strand

(-)primer modification: 5'-dA₂₀HEGL



follow-up treatment: → none
HEGL-spacer as terminator for *Taq* polymerase
(interruption of elongation of the (+)strand)



elongation of the (-)strand by 20 nucleotides